



Moxy[®] 2E

Selective weed control in a variety of crops and non-cropland.


Moxy[®] 2E herbicide controls broadleaf weeds in seedling alfalfa, barley, Conservation Reserve Program (CRP) grasses, corn (field and pop), flax, garlic, grasses grown for seed and sod production, mint, non-cropland/industrial sites, non-residential turf grass, oats, onions (dry bulb), rye, sorghum (grain and forage), triticale and wheat. A contact herbicide, Moxy[®] 2E is most effective when sprayed on actively growing broadleaf weed seedlings.


FEATURES AND BENEFITS


- Labeled for use on black nightshade, blue mustard, bristly starbur, common cocklebur, field pennycress, hairy nightshade, jimsonweed, ladythumb, common lambsquarters, annual sowthistle, tartary buckwheat, common tarweed, wild buckwheat and numerous other broadleaf weed species.
- May be tank mixed with several other herbicides and insecticides.
- Provides a wide window of application in many crops. In corn, it can be applied from emergence up to prior to tassel emergence.



 **REGION**
National

 **USE RATE[†]**
Use rates vary by crop, weed and application timing. See product label for specific application rates

 **APPLICATION GUIDE[†]**
See product label for specific application information

 **ACTIVE INGREDIENTS[†]**
Bromoxynil 33.40%
Inert ingredients 66.60%
Moxy[®] 2E herbicide contains two pounds active ingredient per gallon.

 **PACKAGING**
2 x 2.5-gallon jugs