

Warden[®] CX II

By WINFIELD[®]
UNITED



Seed Care

Insecticide with Fungicides

A seed treatment product for protection against damage from listed insects, seed-borne diseases, and seedling diseases on soybean

ACTIVE INGREDIENTS:

| | |
|----------------------------|---------|
| Thiamethoxam ¹ | 16.30 % |
| Mefenoxam ² | 4.89 % |
| Fludioxonil ³ | 0.82 % |
| Sedaxane ⁴ | 0.82 % |
| Picarbutrazox ⁵ | 0.33 % |

OTHER INGREDIENTS:

| | |
|--------|----------|
| TOTAL: | 100.00 % |
|--------|----------|

¹CAS No. 153719-23-4

²CAS No. 70630-17-0 and CAS No. 69516-34-3

³CAS No. 131341-86-1

⁴CAS No. 874967-67-6

⁵CAS No. 500207-04-5

One gallon of Warden CX II contains 1.52 lb thiamethoxam, 0.46 lb mefenoxam, 0.08 lb fludioxonil, 0.08 lb sedaxane, and 0.03 lb picarbutrazox.

**KEEP OUT OF REACH OF CHILDREN.
CAUTION**

SEE ADDITIONAL PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE ON LABEL.

FIRST AID

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

HOTLINE NUMBER: For 24-Hour Medical Emergency Assistance (Human or Animal) Call **1-877-424-7452** Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call **1-800-424-9300**.

NET CONTENTS: 15 GALLONS

EPA Reg. No. 100-1720-1381

EPA Est. No. 100-NE-001

SCPSR-WIN-1720B-L1 0922

4165413 2/0916/2

Distributed by:

Winfield Solutions, LLC

P. O. Box 64589

St. Paul, MN 55164-0589



(01) 0 08 84464 84647 9

Item# 1725406

11559-L1091245 09-22-22 SCP 01-00 (09-26-22)
Warden CX II – 15 gal. booklet/base – Mech
SCP 11559...

AGI 107035

Print Size – Booklet: 5.75" wide X 6.75" high
Base: 6.75" wide X 6.75" high

Pantone colors: Pantone Black, PMS 255

No. of pages: 12

NOTES

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, polyvinyl chloride (PVC) \geq 14 mils, or Viton™ \geq 14 mils
- Shoes plus socks

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This product is toxic to wildlife, freshwater fish, and estuarine/marine fish and highly toxic to aquatic invertebrates. Runoff may be hazardous to aquatic organisms in neighboring areas. **DO NOT** contaminate water when disposing of equipment washwater or rinsate. Exposed treated seed may be hazardous to wildlife, fish, and aquatic invertebrates. Promptly cover or collect seeds spilled during loading or planting.

Pollinator Precautions

Thiamethoxam is highly toxic to bees exposed to direct treatment, and effects may be possible as a result of exposure to translocated residues in blooming crops.

Groundwater Advisory

Mefenoxam is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Thiamethoxam and fludioxonil have properties and characteristics associated with chemicals detected in groundwater. These chemicals may leach into the groundwater if used in areas where soils are permeable, particularly where the water table is shallow, and may result in groundwater contamination.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having high potential for reaching surface water via runoff for several months or more after application.

continued...

PRECAUTIONARY STATEMENTS *(continued)*

Physical or Chemical Hazards

DO NOT store near or use with oxidizing agents. **DO NOT** mix with or allow to come into contact with oxidizing agents. Hazardous chemical reaction may occur.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER AND USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THE NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the seller, and purchase price will be refunded.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Use is permitted as an end-use seed treatment on agricultural establishments and in commercial seed treatment facilities.

DO NOT use for at-plant applications (e.g., hopper box, planter box). This product is to be applied as a water-based slurry through standard liquid-type seed-treatment equipment. Use seed treaters with atomizers or spinning discs for better product coverage on the seed.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation. Notify state and/or federal authorities and Winfield Solutions, LLC immediately if you observe any adverse environmental effects due to use of this product.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR INSECT AND/OR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.

Treatment of highly mechanically scarred or damaged seed or seed known to be of low vigor and poor quality may result in reduced germination and/or reduction of seed and seedling vigor. Treat a quantity of seed using equipment similar to that planned for treating the total seed lot. Prior to treatment, conduct germination tests on a portion of seed before committing the total seed lot to a selected seed treatment.

Due to seed quality, crop or variety sensitivity, and seed storage conditions beyond the control of Winfield Solutions, LLC, no claims are made to guarantee the germination of seed or propagating material for all crop seed when treated with Warden CX II.

Maximum usage when applying both metalaxyl- and mefenoxam-containing products to the same crop within the same season: DO NOT apply more than the maximum yearly total application rate for the active ingredient as stated on the label of the product containing the lowest yearly total on that crop.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours. Exception: If the seed is treated with the product and the treated seed is planted, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton ≥ 14 mils
- Shoes plus socks

PRODUCT INFORMATION

Warden CX II is a seed treatment product containing the active ingredients thiamethoxam insecticide and fludioxonil, mefenoxam, sedaxane, and picarbutrazox fungicides. Warden CX II protects against damage from listed early-season insects and soil-borne and seed-borne diseases of soybeans.

Thiamethoxam is a systemic seed treatment insecticide belonging to the neonicotinoid class of chemistry. Thiamethoxam protects against listed chewing and sucking insects through contact and ingestion.

Mefenoxam fungicide is active against *Pythium*, *Phytophthora* and systemic downy mildew.

Fludioxonil fungicide is active against *Fusarium* and *Rhizoctonia*, and suppresses seed-borne *Sclerotinia* and *Phomopsis* spp.

Sedaxane fungicide is active against seed decay, seedling blight and damping-off caused by *Rhizoctonia* spp.

Picarbutrazox fungicide is active against diseases caused by *Pythium* and *Phytophthora* spp.

Resistance Management

For resistance management, please note that Warden CX II contains Group 4 (mefenoxam), Group 7 (sedaxane), Group 12 (fludioxonil), and Group U17 (picarbutrazox) fungicides. Any fungal population may contain individuals naturally resistant to Warden CX II and other Group 4, Group 7, Group 12, or Group U17 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

Mefenoxam belongs to the phenylamide class of chemistry, which interferes with fungal RNA synthesis. Sedaxane is a succinate dehydrogenase inhibitor (SDHI) and belongs to the carboxamide class of chemistry, which disrupts cellular respiration and energy generation. Fludioxonil belongs to the phenylpyrrole class of chemistry, which interferes with osmotic signal transduction. Picarbutrazox belongs to the tetra-zolyloximes class of fungicides and is not known to be cross-resistant with fungicides that have different modes of action.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of Warden CX II or other Group 4, Group 7, Group 12, or Group U17 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide application. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or Integrated Pest Management (IPM) recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Winfield Solutions, LLC at 1-855-494-6343. You can also contact your pesticide distributor or university extension specialist to report resistance.

Winfield Solutions, LLC encourages responsible product stewardship to ensure effective long-term control of the fungal diseases on this label.

Warden CX II contains a Group 4A insecticide (thiamethoxam, a systemic insecticide belonging to the neonicotinoid class of chemistry, which includes nicotinic acetylcholine receptor (nAChR) agonists). Insect biotypes with acquired or inherent resistance to Group 4A insecticides may eventually dominate the insect population if Group 4A insecticides are used repeatedly as the predominant method of control for targeted species. This may result in partial or total loss of control of those species by Warden CX II or other Group 4A insecticides.

In order to maintain susceptibility to this class of chemistry:

- Avoid using Group 4A insecticides exclusively for season-long control of insect species with more than one generation per crop season.
- For insect species with successive or overlapping generations, apply Warden CX II or other Group 4A insecticides using a "treatment window" approach. A treatment window is a period of time as defined by the stage of crop development and/or the biology of the pests of concern. Within the treatment window, depending on the length of residual activity, single or consecutive applications may be made using seed, in-furrow, or foliar treatments unless otherwise excluded by product labels. **DO NOT** exceed the maximum amount of this insecticide allowed per growing season.
- Following a treatment window of Group 4A insecticides, rotate to a treatment window of effective products with a different mode of action before making additional applications of Group 4A insecticides.
- A treatment window rotation, along with other IPM practices for the crop and use area, is considered an effective strategy for preventing or delaying a pest's ability to develop resistance to this class of chemistry.
- Use products at their full, specified doses.

- Use appropriate, well-maintained equipment. Use specified water volumes and apply at optimal temperatures in order to obtain optimal treatment.
- When rate ranges are given, use the higher rate within the listed rate range when insect pressure is expected to be high.
- If resistance is suspected, **DO NOT** reapply Warden CX II or any other Group 4A insecticides.

Other Insect Resistance Management (IRM) practices include:

- Incorporating IPM techniques into your insect control program
- Monitoring treated insect populations for loss of field efficacy
- Using tank-mixtures or premixes with insecticides from a different target site of action group as long as the involved products are all registered for the same crop outlet and effective rates are applied

For additional information on Insect Resistance Management:

- Contact your local Cooperative Extension Service specialist, certified crop advisor, and/or product manufacturer for additional insect resistance management recommendations.
- Visit the Insecticide Resistance Action Committee (IRAC) website.

Winfield Solutions, LLC encourages responsible product stewardship to ensure effective long-term control of the insects on this label.

APPLICATION DIRECTIONS

Important: Always re-circulate Warden CX II thoroughly before using.

The typical density of Warden CX II is 9.35 pounds per gallon. Consult the manufacturer of the application equipment you plan to use for suitability for this application and for instructions on operation and calibration of the equipment. Follow the manufacturer application instructions for the seed treatment equipment being used.

Apply Warden CX II as a water-based slurry utilizing standard liquid-type-seed treatment equipment that provides uniform seed coverage. Seed treaters with atomizers or spinning discs are highly recommended for better product coverage on the seed. Uneven or incomplete seed coverage may not give the desired level of disease control. Thoroughly mix the specified amount of Warden CX II into the required amount of water or liquid inoculant for the slurry treater and dilution rate to be used.

Certain crops require addition of inoculants when the seed is treated or planted. Warden CX II is compatible with several liquid inoculant products. Consult the maker of the inoculant product and a Winfield Solutions, LLC representative for directions before applying Warden CX II with inoculants.

The total application volume must be sufficient to provide desired level of coverage. Dilution is typically done with water or liquid inoculants. The minimum slurry volume to achieve adequate coverage is 4.0 fluid ounces per 100 pounds of seed assuming, an average seed size of 3,000 seeds per pound (slurry volume will vary depending on seed size). More diluent may be required to obtain complete coverage.

Continuous agitation or mixing of the slurry mixture is necessary to prevent settling out of the solution.

Allow seed to dry before bagging.

USE RESTRICTIONS

- **DO NOT** use for at-plant applications (e.g., hopper box, planter box). This product is to be used in liquid or slurry treaters only.
- **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift.
- **DO NOT** use for feed, food, or oil purposes.
- Store away from feed and foodstuffs.
- **DO NOT** allow children, pets, or livestock to have access to treated seed.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Promptly cover or collect treated seeds spilled during loading or planting.
- Treated seed must be planted into the soil at a depth greater than 1 inch.
- Dispose of all excess treated seed. Leftover treated seed may be double sown around the headland or buried away from water sources in accordance with local requirements.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in the ethanol by-products that are used for agronomic practice.
- **DO NOT** contaminate water bodies when disposing of planting equipment wash waters.
- Dispose of seed packaging in accordance with local requirements.
- Cover crops planted for erosion control or soil improvement, other than those with immediate or 30-day plantback intervals, may be planted sooner than the 120-day interval.
 - o **DO NOT** graze or harvest the crop for food or feed.
- **DO NOT** use at a rate that will result in more than 0.083 lb thiamethoxam per acre (37.8 g ai/A), 0.004 lb fludioxonil per acre (2.0 g ai/A), and 0.0178 lb picarbutrazox per acre (8.1 g ai/A) per calendar year as a seed-treatment application. **DO NOT** apply more than 0.052 lb ai/A (23 g ai/A) per calendar year of mefenoxam- or metalaxyl-containing products as a seed treatment.
- **DO NOT** apply more than 0.004 lb ai sedaxane per acre (2 g ai/A) per calendar year.
- **DO NOT** apply a neonicotinoid insecticide within 45 days of planting seed treated with Warden CX II.
- This seed has been treated with 0.0756 mg thiamethoxam, 0.0226 mg mefenoxam, 0.0038 mg fludioxonil, 0.0038 mg sedaxane, and 0.0015 mg picarbutrazox per seed.
- This treated seed may be planted on the same acres up to 2 times per year.

ROTATIONAL CROP RESTRICTIONS

In the event of a crop failure or harvest of a crop grown from seed treated with Warden CX II, the field may be replanted according to the following schedule of days after the Warden CX II-treated seed was planted:

Plantback Interval Table

| Immediate Plantback Interval | 30-Day Plantback Interval | 120-Day Plantback Interval |
|---|--|----------------------------|
| Corn (field, pop, seed, and sweet) Soybean | Alfalfa Barley <i>Brassica</i> (Cole) leafy vegetables Buckwheat Canola Cotton Cucurbit vegetables Dried shelled peas (chickpea (garbanzo bean), field pea, lentil, and pigeon pea) Dry bulb onions Fruiting vegetables Leafy vegetables Legume vegetables Mint (peppermint and spearmint) Oats Oilseed crops (black mustard seed, borage seed, field mustard seed, Indian rapeseed, peanuts, rapeseed, and safflower seed) Pearl millet Proso millet Rice (dry seeded) Root vegetables Rye Sorghum Strawberry Sunflowers Teosinte Tobacco Triticale Tuberos and corm vegetables Wheat Wild rice | All Other Crops |

SEED CONTAINER LABEL REQUIREMENTS

The Federal Seed Act requires that bags containing treated seeds shall be labeled with the following statements:

- This seed has been treated with thiamethoxam insecticide and fludioxonil, mefenoxam, picarbutrazox, and sedaxane fungicides.
- **DO NOT** use for feed, food, or oil purposes.

In addition, the U.S. Environmental Protection Agency requires the following statements on bags containing seeds treated with Warden CX II:

- Store away from feed and foodstuffs.
- **DO NOT** allow children, pets, or livestock to have access to treated seed.
- Wear long-sleeved shirt; long pants; shoes and socks; and chemical-resistant gloves when handling treated seed.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Promptly cover or collect treated seeds spilled during loading or planting.
- Treated seed must be planted into the soil at a depth greater than 1 inch.
- Dispose of all excess treated seed. Leftover treated seed may be double sown around the headland or buried away from water sources in accordance with local requirements.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in the ethanol by-products that are used for agronomic practice.
- **DO NOT** contaminate water bodies when disposing of planting equipment wash waters.
- Dispose of seed packaging in accordance with local requirements.
- **Groundwater Advisory:** Mefenoxam is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Thiamethoxam and fludioxonil have properties and characteristics associated with chemicals detected in groundwater. These chemicals may leach into the groundwater if used in areas where soils are permeable, particularly where the water table is shallow.
- **Surface Water Advisory:** This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having high potential for reaching surface water via runoff for several months or more after application.
- **Pollinator Precautions:** Thiamethoxam is highly toxic to bees, and effects are possible as a result of exposure to translocated residues in blooming crops.
- In the event of a crop failure or harvest of a crop grown from this treated seed, the field may be replanted immediately to corn (field, pop, seed, or sweet) or soybean. After 30 days, the field may be replanted to: Alfalfa, barley, *Brassica* (Cole) leafy vegetables, buckwheat, canola, cotton, cucurbit vegetables, dried shelled peas (chickpea (garbanzo bean), field pea, lentil, and pigeon pea), dry bulb onions, fruiting vegetables, leafy vegetables, legume vegetables, mint (peppermint and spearmint), oats, oilseed crops (black mustard seed, borage seed, crambe seed, field mustard seed, flax seed, Indian mustard seed, Indian rapeseed, peanuts, rapeseed, and safflower seed), pearl millet, proso millet, rice (dry seeded), root vegetables, rye, sorghum, strawberry, sunflowers, teosinte, tobacco, triticale, tuberous and corm vegetables, wheat, and wild rice. For any other crop, the minimum plant back interval is 120 days from the date this treated seed was planted. A cover crop other than the crops listed above that is planted for erosion control or soil improvement may be planted sooner than the 120-day interval.
 - o **DO NOT** graze or harvest the crop for food or feed.
- **DO NOT** use at a rate that will result in more than 0.083 lb thiamethoxam per acre (37.8 g ai/A), 0.004 lb fludioxonil per acre (2.0 g ai/A), and 0.0178 lb picarbutrazox per acre (8.1 g ai/A) per calendar year as a seed-treatment application. **DO NOT** apply more than 0.052 lb ai/A (23 g ai/A) per calendar year of mefenoxam- or metalaxyl-containing products as a seed treatment.
- Maximum usage when applying metalaxyl- and mefenoxam-containing products to the same crop within the same season: **DO NOT** apply more than the maximum yearly total application rate for the active ingredient as stated on the label of the product containing the lowest yearly total on that crop.
- **DO NOT** apply more than 0.004 lb ai sedaxane per acre (2 g ai/A) per calendar year.

- This seed has been treated with 0.0756 mg thiamethoxam, 0.0226 mg mefenoxam, 0.0038 mg fludioxonil, 0.0038 mg sedaxane, and 0.0015 mg picarbutrazox per seed.
- **DO NOT** apply a neonicotinoid insecticide within 45 days of planting seed treated with Warden CX II.
- This treated seed may be planted on the same acres up to 2 times per year.

CROP USE DIRECTIONS

When applied according to the **USE RATE TABLE**, Warden CX II provides early-season protection against injury by aphids, bean leaf beetle, grape colaspis, leaf miners, leaf hoppers, Mexican bean beetle, seed corn maggot, three-cornered alfalfa hopper, thrips, white grubs, and wireworm.

Warden CX II provides protection against damping-off and seed borne rots due to *Pythium*, *Phytophthora*, *Fusarium*, and *Rhizoctonia* species and early-season Phytophthora root rot. Warden CX II also suppresses seed-borne *Sclerotinia* and *Phomopsis* species.

USE RATE TABLE

| Crop | Rate of Warden CX II | | | |
|---------|---------------------------------------|---|--|--------------------------------|
| Soybean | Fl oz per 100 lb seed ¹ | Fl oz per 140,000 seeds ¹ | Grams ai per 100 kg seed | Mg ai per seed ² |
| | 4.2 | 1.96 | Thiamethoxam: 50 Mefenoxam: 15.0 Fludioxonil: 2.5 Sedaxane: 2.5 Picarbutrazox: 1.0 | Total = 0.1074 mg |

¹ The fl oz Warden CX II per 100 lb seed and per 140,000 seed rates are based on 3,000 seeds per pound.

² This seed has been treated with 0.0756 mg ai/seed of thiamethoxam, 0.0226 mg ai/seed of mefenoxam, 0.0038 mg ai/seed of fludioxonil, 0.0038 mg ai/seed of sedaxane, and 0.0015 mg ai/seed picarbutrazox.

Stored Grain Protection

When treated according to the directions for post-planting protection against listed pests, Warden CX II will also provide protection during post treatment storage of the soybean seed against damage from the following stored grain insects: Indian Meal Moth (*Plodia interpunctella*), Rice Weevil (*Sitophilus oryza*), Red Flour Beetle (*Tribolium castaneum*), and Lesser Grain Borer (*Rhizopertha dominica*).

If the soybean seed to be treated has existing infestations of stored grain insects, fumigate the seed with a registered product approved for such use, prior to treating with Warden CX II and bagging.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store in a cool, dry place. Storage for extended periods above 90°F is not recommended.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

Container Handling (5 gallons or less)

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Container Handling (greater than 5 gallons)

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other side and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Container Handling (greater than 5 gallons)

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or re-circulate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

Warden® is a registered trademark of Winfield Solutions, LLC

Viton™ is a trademark of The Chemours Company FC, LLC

©2022 Winfield Solutions, LLC

FOR CHEMICAL EMERGENCY: Spill, leak, fire, or accident, call CHEMTREC 1-800-424-9300

Distributed by:
Winfield Solutions, LLC
P. O. Box 64589
St. Paul, MN 55164-0589

Warden® CX II

By **WINFIELD®
UNITED**

Insecticide with Fungicides

A seed treatment product for protection against damage from listed insects, seed-borne diseases, and seedling diseases on soybean

| | |
|---------------------------------|----------|
| Active Ingredients: | |
| Thiamethoxam ¹ | 16.30 % |
| Mefenoxam ² | 4.89 % |
| Fludioxonil ³ | 0.82 % |
| Sedaxane ⁴ | 0.82 % |
| Picarbtrazox ⁵ | 0.33 % |
| Other Ingredients: | 76.84 % |
| Total: | 100.00 % |

¹CAS No. 153719-23-4

²CAS No. 70630-17-0 and CAS No. 69516-34-3

³CAS No. 131341-86-1

⁴CAS No. 874967-67-6

⁵CAS No. 500207-04-5

One gallon of Warden CX II contains 1.52 lb thiamethoxam, 0.46 lb mefenoxam, 0.08 lb fludioxonil, 0.08 lb sedaxane, and 0.03 lb picarbtrazox.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

EPA Reg. No. 100-1720-1381

EPA Est. No. 100-NE-001

SCSPS-WIN-1720B-L1 0922
4165413
2/0916/2

NET CONTENTS: 15 GALLONS

Distributed by:
Winfield Solutions, LLC
P. O. Box 64589
St. Paul, MN 55164-0589

KEEP OUT OF REACH OF CHILDREN. CAUTION

See additional precautionary statements and directions for use on label.

FIRST AID

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

HOTLINE NUMBER: For 24-Hour Medical Emergency Assistance (Human or Animal) Call **1-877-424-7452** Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call **1-800-424-9300**.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

Environmental Hazards: This product is toxic to wildlife, freshwater fish, and estuarine/marine fish and highly toxic to aquatic invertebrates. Runoff may be hazardous to aquatic organisms in neighboring areas. **DO NOT** contaminate water when disposing of equipment washwater or rinsate. Exposed treated seed may be hazardous to wildlife, fish, and aquatic invertebrates. Promptly cover or collect seeds spilled during loading or planting.

Pollinator Precautions: Thiamethoxam is highly toxic to bees exposed to direct treatment, and effects may be possible as a result of exposure to translocated residues in blooming crops.

Groundwater Advisory: Mefenoxam is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Thiamethoxam and fludioxonil have properties and characteristics associated with chemicals detected in groundwater. These chemicals may leach into the groundwater if used in areas where soils are permeable, particularly where the water table is shallow, and may result in groundwater contamination.

Surface Water Advisory: This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground-

| | | | |
|---------------|-------|-----|-------------|
| THIAMETHOXAM | GROUP | 4A | INSECTICIDE |
| MEFENOXAM | GROUP | 4 | FUNGICIDE |
| FLUDIOXONIL | GROUP | 12 | FUNGICIDE |
| SEDAXANE | GROUP | 7 | FUNGICIDE |
| PICARBUTRAZOX | GROUP | U17 | FUNGICIDE |

water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application.

Physical or Chemical Hazards: DO NOT store near or use with oxidizing agents. **DO NOT** mix with or allow to come into contact with oxidizing agents. Hazardous chemical reaction may occur.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place. Storage for extended periods above 90°F is not recommended.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

Container Handling

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other side and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

Warden® is a registered trademark of Winfield Solutions, LLC

©2022 Winfield Solutions, LLC

FOR CHEMICAL EMERGENCY: Spill, leak, fire, or accident, call CHEMTREC 1-800-424-9300