

# Stonewall® 540


By **WINFIELD UNITED**

**ACTIVE** Glyphosate

Stonewall® 540 provides grass and broadleaf weed control in glyphosate tolerant crops.

## FEATURES & BENEFITS

- Stonewall 540 is a post-emergent, non-selective herbicide that is absorbed and translocated through weeds for “shoots-to-roots” control.
- Stonewall 540 controls more than 100 annual and perennial weeds and includes versatile use patterns including in-crop applications in Roundup Ready® and TruFlex™ canola crops, as well as pre-seed, pre-harvest and post-harvest applications.
- Hard water can reduce the activity of glyphosate. Test your water and add Crimson® NG to reduce the effects of hard water and other cations.



### FOR USE ON

Glyphosate tolerant: corn, canola, soybean and sugarbeet. Plus pre-seed, pre-harvest, post-harvest, chemfallow and vegetation management.

For pre-harvest applications visit [www.keepingitclean.ca](http://www.keepingitclean.ca) and observe appropriate pre-harvest intervals.

### Application Timing

Varies by crop type, see label for more information.

### WEEDS CONTROLLED

Ensure appropriate rate is selected based on weed type and weed stage.

Alfalfa, annual sow thistle, barnyard grass, Canada fleabane, Canada thistle, chickweed, cleavers, corn spurry, cow cockle, crabgrass, curled dock, dandelion, downy brome, field bindweed, flixweed, foxtail barley, green foxtail, green smartweed, hemp-nettle, Japanese knotweed, kochia (non-glyphosate resistant biotypes), lady's-thumb, lamb's-quarters, narrow-leaved hawk's beard, night flowering catchfly, perennial sow thistle, Persian dandelion, prickly lettuce, quackgrass, ragweed, redroot pigweed, round-leaved mallow, Russian thistle, shepherd's purse, smooth brome, smooth pigweed, stinkweed, stork's-bill, velvetleaf, volunteer canola, volunteer cereals, volunteer flax, volunteer mustard, wild mustard, wild buckwheat, wormwood.

### Application Rates

0.2-3.2 L/acre. Select rate based on weed species and stage based on registered use pattern.

### Adjuvant Rates

- Crimson NG up to 2.5% v/v
- InterLock® for drift control

### Minimum Water Volume

- Ground application 5 US gal/acre
- Aerial application 2 US gal/acre\*

### Rainfast

- 1 hour

\*Do not apply by air except for pre-harvest aerial application.



### COMMON USE RATE

Varies based on crop and weed stage



### PACKAGING

2 x 10 L jugs/case  
115 L drum  
500 L, 800 L, 1000 L tote



### ACTIVE INGREDIENT

Glyphosate (K+) 540 g/L



### HERBICIDE GROUP

Group 9

Always tank mix with multiple modes of action targeted at the most troublesome weeds or those most prone to herbicide resistance.